

Intranight optical variability of Blazars

Ram Sagar¹, C. S. Stalin^{1,2,3}, Gopal-Krishna² and Paul J. Wiita⁴

¹ *State Observatory, Manora Peak, Nainital 263 129, India*

² *National Centre for Radio Astrophysics, TIFR, Pune University Campus, Post Bag 3, Pune 411 007, India*

³ *Laboratoire de Physique Corpusculaire et Cosmologie, Collège de France, 11 pl. Marcelin Berthelot, F-75231, Paris Cedex 5, France*

⁴ *Department of Physics & Astronomy, MSC 8R0314, Georgia State University, Atlanta, Georgia 30303-3088, USA*

Accepted 2003 October 23, Received 2003 October 3; in original form 2003 May 21

ABSTRACT

We present results of a multi-epoch intra-night optical monitoring of eleven blazars consisting of six BL Lac objects and five radio core-dominated quasars (CDQs). These densely sampled and sensitive R-band CCD observations, carried out from November 1998 through May 2002 during a total of 47 nights with an average of 6.5 hours/night, have enabled us to detect variability amplitudes as low as $\sim 1\%$ on intra-night time scales. A distinction is found for the first time between the intra-night optical variability (INOV) properties of these two classes of relativistically beamed radio-loud AGNs. BL Lacs are found to show a duty cycle (DC) of INOV of $\sim 60\%$, in contrast to CDQs, which show a much smaller INOV DC of $\sim 20\%$, the difference being attributable mainly to the weakly polarized CDQs. On longer time scales (i.e., between a week to a few years) variability is seen from all the CDQs and BL Lacs in our sample. The results reported here form part of our long-term programme to understand the intra-night optical variability characteristics of the four main classes of luminous AGNs, i.e., radio-quiet quasars (RQQs) and radio lobe-dominated quasars (LDQs), as well as CDQs and BL Lac objects.

Key words: galaxies: active – galaxies: jets – galaxies: photometry – quasars: general

1 INTRODUCTION

Variability observations of active galactic nuclei (AGNs) on intra-night time scales can provide valuable clues to the physics of the innermost nuclear regions in these objects. Blazars (core-dominated quasars: CDQs, and BL Lac objects: BL Lacs) as a class of AGNs are characterised by the most violent variations at almost all wavelengths over a wide range of time scales. Blazars' properties are consistent with relativistic beaming, that is bulk relativistic motion of the jet plasma at small angles to the line of sight, which gives rise to strong amplification and rapid variability in the observer's frame. CDQs and BL Lacs are thought to be the beamed counterparts of high and low luminosity radio galaxies, respectively (e.g., Urry & Padovani 1995). The main difference between CDQs and BL Lacs lies in their emission lines, which are strong in CDQs, but weak, and in many cases, undetected, in BL Lacs. CDQ spectra can extend up to around GeV energies, whereas the spectra of BL Lacs can extend up to TeV energies. Though it is very likely that both CDQs and BL Lacs are dominated by non-thermal Doppler boosted jets, some important differences have been found between their apparent non-thermal properties, such as the magnetic field patterns in their parsec-scale jets (Gabuzda et al. 1992; but see Gopal-Krishna & Wiita 1993).

The intra-night optical variability (INOV), or microvariability, of blazars has been an established phenomenon for over a dozen years (Miller, Carini & Goodrich 1989; Carini et al. 1991). Although the origin(s) of INOV in all AGNs is still uncertain, for blazars it is generally associated with the non-thermal Doppler boosted emission from jets (Blandford & Rees 1978; Marscher & Gear 1985; Marscher, Gear & Travis 1992; Hughes, Aller & Aller 1992; Wagner & Witzel 1995). Still, alternative models, which invoke accretion disk instabilities or perturbations (e.g., Mangalam & Wiita 1993; for a review, see Wiita 1996) may also explain some INOV, particularly in radio-quiet quasars (RQQs) where any contribution from the jets, if they are at all present, is weak. Several studies of the INOV of blazars are available in the literature (e.g., Heidt & Wagner 1996; Dai et al. 2001; Romero et al. 2002; Xie et al. 2002). However, the unique feature of the present study is the deliberate focus on the comparison of the INOV properties of the two Doppler beamed AGN classes, namely CDQs and BL Lacs. Other results from this large programme, involving the nature of INOV in BL Lacs and RQQs (Gopal-Krishna et al. 2003, hereafter GSSW03), and a comparison of INOV between lobe-dominated radio-loud quasars (LDQs) and RQQs (Stalin et al. 2003a), have been published elsewhere.

Table 1. The sample of core-dominated quasars and BL Lacs monitored in the present programme

Object	Other Name	Type	RA(2000)	Dec(2000)	B (mag)	M_B (mag)	z	α	%Pol* (opt)	$R^\dagger$
0219+428	3C 66A	BL	02 22 39.6	+43 02 08	15.71	-26.5	0.444	-0.19§	11.70	676.1
0235+164	AO 0235+164	BL	02 38 38.9	+16 37 00	16.46	-27.6	0.940	0.67	14.90	1949.8
0735+178	PKS 0735+17	BL	07 38 07.4	+17 42 19	16.76	-25.4	>0.424	-0.26§	14.10	3548.1
0851+202	OJ 287	BL	08 54 48.8	+20 06 30	15.91	-25.5	0.306	0.18§	12.50	2089.3
0955+326	3C 232	CDQ	09 58 20.9	+32 24 02	15.88	-26.7	0.530	-0.11§	0.53	549.5
1128+315	B2 1128+31	CDQ	11 31 09.4	+31 14 07	16.00	-25.3	0.289	-0.41	0.62	269.2
1215+303	B2 1215+30	BL	12 17 52.0	+30 07 01	16.07	-24.8	0.237	-0.17§	8.00	426.6
1216-010	PKS 1216-010	CDQ	12 18 35.0	-01 19 54	16.17	-25.9	0.415	-0.03§	6.90	218.8
1225+317	B2 1225+31	CDQ	12 28 24.8	+31 28 38	16.15	-30.0	2.219	0.01	0.16	182.0
1308+326	B2 1308+32	BL	13 10 28.7	+32 20 44	15.61	-28.6	0.997	-0.09§	10.20	512.9
1309+355	PG 1309+355	CDQ	13 12 17.7	+35 15 23	15.60	-24.7	0.184	-0.12	0.31	22.9

* Reference for optical polarizations: Wills et al. (1992)

 † R is the ratio of the radio-to-optical flux densities as defined in the text

§ Reference for these radio fluxes: Kovalev et al. (1999); for others see text

2 SAMPLE, OBSERVATIONS AND REDUCTIONS

The sample of blazars used in this work consists of 6 BL Lac objects and 5 CDQs. All are bright, with apparent B magnitudes between 15.6 and 16.8, so that short exposures can still provide good signal to noise ratios. At most two of these, one CDQ (1225+317, with $z = 2.219$) and possibly one BL Lac (0735+178, with $z > 0.424$), lie at $z > 1$, so this sample provides a fairly even coverage of the redshift range up to $z = 1$ for each blazar subclass. Note that we have adopted a BL Lac classification for B2 1308+326, following the 1-Jy (Stickel, Fried & Kühr 1993) and Padovani & Giommi (1995) catalogs, even though it is classified as a CDQ in the Véron-Cetty & Véron (1998) catalog. Also, for the BL Lac object PKS 0735+178 we have adopted a redshift $z = 0.424$, though formally the published value is $z > 0.424$ (see Véron-Cetty & Véron 1998). For each object we could find a measurement of the degree of optical polarization (Wills et al. 1992); it can be seen from Table 1 that for four out of the five CDQs the percentage polarization was $< 1\%$ at the time those measurements were made, and hence they can be assigned to the sub-class of low-polarization CDQs.

Basic information of these objects are given in Table 1. The values of the radio spectral index, α (with $S_\nu \propto \nu^\alpha$), given in Table 1 were usually determined from linear spectral fitting to the available *near simultaneous* flux density measurements between 1 and 22 GHz reported by Kovalev et al. (1999); those not appended with § are based on linear fits to *non-simultaneous* flux measurements taken from NED¹.

The observations were made using the 104-cm Sampurnanand telescope of the State Observatory, Naini Tal which is an RC system with a f/13 beam (Sagar 1999). The detectors used were a cryogenically cooled 1024×1024 CCD chip (prior to October 1999) and a 2048×2048 chip (after October 1999), both mounted at the Cassegrain focus. Each pixel of both the CCDs correspond to a square of 0.38 arcsec on the sky, covering a total field of $\sim 12' \times 12'$ in the case of the larger CCD and $\sim 6' \times 6'$ in the case of the

smaller CCD. Observations were almost always done using an R filter, as it was near the maximum response of the CCD system and thus allowed us to achieve good temporal resolution; however, on two nights quasi-simultaneous observations were done using R and I filters. To improve S/N, observations were carried out in 2×2 binned mode. On each night only one QSO was monitored as continuously as possible and the typical sampling rate was about 5 frames per hour. The choice of the exposure time depended on the brightness state of the QSO, the moon's phase and sky transparency. The field containing the QSO was adjusted so as to have at least 2 (and usually 3) comparison stars within about a magnitude of the QSO on the CCD frame.

Preliminary processing of the images as well as photometry was done using the IRAF² software. Photometry of the QSO and the comparison stars recorded on the same CCD frame was carried out using the *phot* task in IRAF. The same circular aperture was used for the photometry of the QSO and the comparison stars for all the images acquired over the night. This optimum aperture was selected by considering a range of apertures starting from the median FWHM over the night for the photometry and choosing that aperture that produced the minimum variance in the star - star differential lightcurve (DLC) of the steadiest pair of comparison stars. Further details of the observations and reductions are presented elsewhere (Stalin 2002; Stalin et al. 2003b). DLCs of the AGN relative to the comparison stars as well as between all pairs of comparison stars (usually three, but in some cases, two) are constructed from the derived instrumental magnitudes. The DLCs of the AGN relative to the comparison stars are used to look for the presence of INOV in the AGN. The choice of more than one comparison star in the differential photometry enables us to reliably identify QSO variability, as any stars which themselves varied during the night can be identified and discarded. The position and apparent magnitudes of the comparison stars used in the differential photometry of our sample of blazars from the USNO catalog³ are given in Table 2. Note that the magni-

² Image Reduction and Analysis Facility, distributed by NOAO, operated by AURA, Inc. under agreement with the US NSF.

³ <http://archive.eso.org/skycat/servers/usnoa>

¹ URL <http://ned.ipac.caltech.edu/>






Figure 1. Differential R-band light curves for the BL Lacs and core-dominated quasars. The name of the object, the date and duration of observation are given on the top of each panel. The upper panel(s) give the DLC of the pair(s) of comparison stars whereas the subsequent lower panels are the DLCs of the quasar relative to the comparison stars. Note the different scales for many of the sources. I-band DLCs are shown as open symbols for two nights for BL 0219+428.

Figure 1. *Continued*


 Figure 1. *Continued*

tudes of the comparison stars taken from this catalog have uncertainties of up to 0.25-mag.

3 RESULTS

3.1 Differential light curves (DLCs)

DLCs are presented for those AGNs which have shown clear evidence of INOV in Fig. 1. We consider a source to be variable only if correlated variations (both in amplitude and time) are found in the DLCs of the AGN relative to all the comparison stars considered. All of the six BL Lacs in our sample showed INOV on at least one night, whereas this was

Table 2. Positions and apparent magnitudes of the comparison stars

Source	Star	RA(2000) h m s	Dec(2000) d m s	R mag	B mag
0219+428 BL	S1	02 22 45.13	43 04 19.6	14.2	15.6
	S2	02 22 47.23	43 06 00.1	13.9	14.7
	S3	02 22 28.39	43 03 40.7	13.9	14.8
0235+164 BL	S1	02 38 56.44	16 38 56.5	14.5	15.7
	S2	02 38 54.48	16 36 03.1	15.5	16.1
	S3	02 38 38.53	16 40 05.2	16.7	18.0
0735+178 BL	S1	07 38 03.45	17 42 56.1	16.4	16.5
	S2	07 38 17.10	17 39 03.7	16.0	16.0
	S3	07 38 10.29	17 43 43.9	16.4	16.6
0851+202 BL	S1	08 54 46.11	20 07 20.3	15.7	16.8
	S3	08 54 43.70	20 02 42.2	16.4	18.8
0955+326 CDQ	S1	09 58 14.45	32 23 45.8	14.0	15.0
	S2	09 58 18.32	32 28 35.1	14.5	14.9
	S3	09 58 26.47	32 26 54.3	15.9	16.5
1128+310 CDQ	S1	11 31 02.12	31 11 39.3	15.9	16.7
	S2	11 30 54.41	31 11 47.8	15.6	15.8
	S3	11 31 18.04	31 17 16.8	15.5	17.1
1215+303 BL	S1	12 17 45.96	30 04 51.0	14.4	16.8
	S2	12 17 49.13	30 07 02.3	15.8	16.4
	S3	12 17 44.47	30 09 44.1	13.7	14.5
	S4	12 18 09.03	30 09 35.8	14.7	15.6
	S5	12 17 26.62	30 07 53.5	14.2	15.3
1216-010 CDQ	S1	12 18 42.91	-01 19 24.8	15.4	16.2
	S2	12 18 45.07	-01 19 47.2	15.3	15.8
1225+317 CDQ	S1	12 28 18.78	31 25 20.1	14.6	15.5
	S2	12 28 30.62	31 26 34.2	15.6	16.6
	S3	12 28 13.60	31 27 36.3	17.1	18.4
	S4	12 28 29.15	31 25 18.5	15.5	16.7
1308+326 BL	S1	13 10 19.69	32 23 53.6	16.9	17.8
	S2	13 10 18.09	32 20 07.3	15.8	16.6
	S3	13 10 29.69	32 25 54.0	16.4	16.8
	S4	13 10 38.31	32 17 31.7	16.0	16.7
1309+355 CDQ	S1	13 12 39.17	35 19 50.4	14.6	16.0
	S2	13 12 40.49	35 16 03.0	15.0	15.9
	S3	13 12 30.99	35 16 08.7	14.3	15.8

the case for only two of the five CDQs monitored (Table 3). For each variable AGN, we have statistically quantified the variability and have derived variability parameter, amplitude and time scale of variability below.

3.1.1 Variability parameter (C_{eff})

To quantify the variability, we have employed a statistical criterion based on the parameter C , similar to that followed by Jang & Miller (1997), with the added advantage that for each AGN we have DLCs relative to multiple comparison stars. This allows us to discard any variability candidates for which the multiple DLCs do not show clearly correlated trends, both in amplitude and time. We define C for a given DLC as the ratio of its standard deviation, σ_T , and the mean σ of its individual data points, $\eta\sigma_{\text{err}}$. Here η is the factor by which the average of the measurement errors (σ_{err} , as given by *phot*) should be multiplied. It has been argued in the literature that the final errors given by DAOPHOT/IRAF are often too small (Gopal-Krishna et al. 1995; Garcia et al. 1999). We find $\eta = 1.50$ (Stalin 2002; GSSW03). The value of C_i for the i^{th} DLC of the AGN has the correspond-

ing probability, p_i , that the DLC is steady (non-variable), assuming a normal distribution. For a given AGN we then compute the joint probability, P , by multiplying the values of p_i 's for individual DLCs available for the AGN. The effective C parameter, C_{eff} , corresponding to P , is given in Table 3 for each variable AGN (i.e., $C_{\text{eff}} > 2.57$, corresponds to a confidence level of variability in excess of 99%). We also note that for these AGN all the DLCs involving only comparison stars were found to show statistically insignificant variability. This is quantified by giving within parentheses the values of C_{eff} for the DLC involving two stable comparison stars.

3.1.2 Amplitude of variability (ψ)

For objects which are variable we define the variability amplitude as (Romero, Cellone & Combi 1999)

$$\psi = \sqrt{(D_{\text{max}} - D_{\text{min}})^2 - 2\sigma^2}, \quad (1)$$

with

$$\begin{aligned} D_{\text{max}} &= \text{maximum in the quasar differential light curve} \\ D_{\text{min}} &= \text{minimum in the quasar differential light curve} \\ \sigma^2 &= \eta^2 \langle \sigma_{\text{err}}^2 \rangle. \end{aligned}$$

The variability amplitudes computed using Eq. (1) for objects which have shown INOV are given in Table 3 in per cent. Note that the smallest clearly detected amplitude of INOV in the present sample is 1%.

3.1.3 Structure function

The structure function is frequently used to characterize variability properties such as time-scales and periodicities present in light curves. The first order structure function for a DLC containing N evenly spaced data points is defined as (Simonetti, Cordes & Heeschen 1985)

$$D_X^1(\tau) = \frac{1}{N(\tau)} \sum_{i=1}^N w(i)w(i+\tau)[X(i+\tau) - X(i)]^2 \quad (2)$$

where τ = time lag, $N(\tau) = \sum w(i)w(i+\tau)$ and the weighting factor $w(i)$ is 1 if a measurement exists for the i^{th} interval, 0 otherwise. The error in each point in the computed structure function is

$$\sigma^2(\tau) = \frac{8\sigma_{\delta X}^2}{N(\tau)} D_X^1(\tau) \quad (3)$$

where $\sigma_{\delta X}^2$ is the measurement noise variance.

Since the samplings of our DLCs are quasi-uniform, we have determined structure functions using an interpolation algorithm. For any time lag τ , the value of $X(i+\tau)$ was calculated by linear interpolation between the two adjacent data points. A typical time scale in the light curve (i.e., the time between a maximum and a minimum, or vice versa) is indicated by a local maximum in the structure function. In case of a monotonically increasing structure function, the source possesses no typical time-scale shorter than the total duration of observations. The plots of the structure function for objects which have shown definite micro-variability are given in Fig. 2, and the inferred time scales of variabilities are given in Table 3. The behaviour exhibited by these SF plots for our observations are the following types:

(i) Sources whose SF display no plateau even at long time lag. The interpretation is that any characteristic time-scale, τ , is longer than the duration of the light curve and therefore the longest time lag available represents only a lower limit to any characteristic time-scale of variation;

(ii) Sources whose SF show two plateaux. This indicates the presence of two time-scales, which may possibly be related to different physical processes.

(iii) Sources which exhibit one plateau followed by a dip in the structure function. The plateau is interpreted as the variability time-scale and the dip as a period of a possibly cyclic signal in the light curve; these possible periods are denoted by P in Table 3.

3.2 Long term optical variability (LTOV)

Our observations also provide information on the LTOV. The number of epochs covered range between three and seven. For the six BL Lacs in our sample the total time-spans covered range between about two to three years. Similarly, year-like time coverage is available for four of our five CDQs; for the CDQ 1216–010 the overall time-spans are much shorter (~ 1 week). Interestingly, even in this case, LTOV is convincingly detected just as in the case of the remaining CDQs and BL Lacs in our sample. The smallest amplitude of LTOV was found in the CDQ 1225+317; a $\sim 2\%$ brightening within a year between April 2000 and April 2001. It is noteworthy that this CDQ has not only the highest redshift ($z = 2.219$) but also the lowest optical polarization ($P_{\text{opt}} = 0.16\%$) in our sample. Also a $\sim 2\%$ brightening is noticed in 1216–010 within 48 hours.

The LTOV results are summarized in the last column of Table 3, where the differences in the nightly means from those of the previous observations are tabulated. Consequently, for an object, the last column is left blank at the first epoch of its observations. We also give comments on the LTOV of individual sources below.

4 NOTES ON THE VARIABILITY OF INDIVIDUAL SOURCES

BL 0219+428: This is the best observed blazar in our sample, with seven epochs of monitoring with durations between five and ten hours, over the period from November 1998 to December 2000. Fig. 1 shows the DLCs for the six of these epochs when INOV was seen. On two of these epochs in October 2000, we also have I band monitoring in addition to the default R band monitoring. On both of those nights the INOV in I and R bands is found to be strongly correlated. The overall INOV amplitudes on the six nights range between 2 and 8%. On 13 November 1999 and 1 December 2000, fairly abrupt changes in the slope of the DLCs were seen during the 5 to 6 hours of continuous monitoring, and the INOV amplitudes recorded on these two nights are also the largest observed for this object (Table 3). Another remarkable feature is that on 13 November 1999 at 19.6 UT a 1.5% downward “glitch” occurred within a time span of less than 10 minutes, followed by the onset of steady fading. An expanded version of this part of the DLC is shown in the top panel of column 1 in Fig. 1. The SF plots for the six

Table 3. Log of INOV and LTOV observations of CDQs and BL Lacs. In the case of INOV, the variability parameter (C_{eff}), amplitude (ψ), characteristic time scale (τ) and Period (P) of variability are also given. The number within parentheses in the C_{eff} column denotes the value for the DLC involving two stable comparison stars.

Object	Date	No. of points	Duration (hours)	INOV status*	C_{eff} AGN(star)	ψ (%)	τ (hours)	P (hours)	LTOV (Δ m)
0219+428 BL	14.11.98	118	6.5	V	6.0 (0.5)	5.4	> 6.5		
	13.11.99	123	5.7	V	> 6.6(1.2)	5.5	> 5.9		+0.25
	24.10.00	73	9.1	V	5.8 (1.5)	4.3	> 9.1		+0.45
	26.10.00	82	10.1	V	3.5 (1.4)	3.2	4.9		-0.26
	01.11.00	103	9.0	V	2.9 (1.1)	2.2	3.9		-0.08
	24.11.00	71	5.1	NV					+0.06
	01.12.00	59	5.1	V	>6.6 (1.1)	8.0	> 5.1		-0.35
0235+164 BL	13.11.98	36	4.4	—	—	—	—	—	
	12.11.99	39	6.6	V	>6.6 (1.1)	12.8	3.6		-0.67
	14.11.99	34	6.2	V	3.2 (1.1)	10.3	3.4		+0.83
	22.10.00	39	7.9	V	2.6 (1.4)	7.6			+0.45
	28.10.00	29	6.8	—	—	—	—	—	+0.35
0735+178 BL	26.12.98	49	7.8	NV					
	30.12.99	65	7.4	NV					-0.45
	25.12.00	43	6.0	NV					-0.70
	24.12.01	43	7.3	V	2.8 (2.0)	1.0	> 8.1		+0.43
0851+202 BL	29.12.98	19	6.8	V	2.8 (0.4)	2.3	> 6.8		
	31.12.99	29	5.6	V	6.5 (1.1)	3.8	3.0		+0.60
	28.03.00	22	4.2	V	5.8 (0.8)	5.0	1.2		-0.93
	17.02.01	48	6.9	V	2.7 (0.7)	2.8	2.0	3.8	-0.70
0955+326 CDQ	19.02.99	36	6.5	NV					
	03.03.00	37	6.3	NV					-0.05
	05.03.00	34	6.9	PV	2.2 (0.6)	0.7			0.00
1128+315 CDQ	18.01.01	31	5.7	NV					
	09.03.02	27	8.2	NV					-0.14
	10.03.02	28	8.3	NV					0.00
1215+303 BL	20.03.99	21	7.0	V	5.5 (0.9)	3.5	4.2		
	25.02.00	28	5.9	NV					+0.19
	31.03.00	27	5.0	NV					+0.13
	25.04.01	29	6.5	—	—	—	—	—	+0.21
	19.04.02	23	6.8	V	4.9 (1.3)	1.8	> 6.8		-0.11
1216+010 CDQ	11.03.02	22	8.0	V	3.2 (0.9)	7.3	1.8	3.2	
	13.03.02	24	8.5	V	2.6 (1.5)	3.8	1.2	2.2	-0.02
	15.03.02	11	3.9	V	3.9 (0.9)	5.5	1.0	2.2	+0.11
	16.03.02	22	8.2	V	6.6 (1.4)	14.1	> 8.2		-0.14
1225+317 CDQ	07.03.99	49	6.6	NV					
	07.04.00	23	6.0	NV					0.00
	20.04.01	34	7.4	NV					-0.02
1308+326 BL	23.03.99	17	6.0	—	—	—	—	—	
	26.04.00	16	5.6	NV					+0.21
	03.05.00	19	6.7	—	—	—	—	—	0.00
	17.03.02	19	7.7	V	3.1 (0.6)	3.4	1.2,4.4		-1.90
	20.04.02	14	5.8	NV					+0.44
	02.05.02	15	5.1	NV					-0.03
1309+355 CDQ	25.03.99	39	6.7	NV					
	01.04.01	32	4.6	NV					+0.10
	02.04.01	41	5.2	NV					0.00

* V = variable, NV = not variable, PV = probably variable

nights are smoother than usual, although significant flattenings are apparent on the nights of 01 November 2000 and 26 October 2000, indicating time scales of around 4 hours (Fig. 2). Both brightening and fading were noted in this BL Lac's LTOV. The blazar faded by 0.70 mag during our first three epochs of observations (between 14 November 1998 and 24 October 2000). This was followed by a brightening of 0.34 mag within a week that encompassed two additional nights of study. The object had dimmed by 0.06 mag when observed 23 days later on 24 November 2000 and was found

to have brightened by 0.35 mag when observed for the last time on 1 December 2000.

BL 0235+164: This well known BL Lac object was monitored on five epochs between November 1998 and October 2000. The DLCs of the first and last epochs are quite noisy, owing to the less sensitive CCD chip in the former case and faintness of the object in the latter case. On each of the remaining three DLCs, INOV is clearly seen against all three comparison stars, with amplitudes ranging between 7.6% and 12.8% (Fig. 1). The derived SFs for the two nights in





Figure 2. First order structure function of BL Lacs and CDQs which have shown INOV on the marked epoch against the indicated comparison star.

November 1999 have good S/N and in both cases a flattening is noticed corresponding to time scale of around 3 hours (Fig. 2). INOV monitoring of this object has also been recently reported by Xie et al. (2001). A particularly interesting feature of the DLCs on 12 November 1999 is the rapid brightening of this object by about 2% within about 22 minutes (which corresponds to only about 11 minutes in the rest frame). This BL Lac also showed significant LTOV, with changes in both directions during the slightly under two year period during which we observed it. It had initially brightened by 0.67 mag between the first two epochs of observations (13

November 1998 and 12 November 1999) followed by a dramatic 0.83 mag fading in two days (12 November 1999 to 14 November 1999). Additional fading by 0.45 mag was noticed when it was observed a year later on 22 October 2000. This was again followed by a fading of 0.35 mag in the following week (between 22 October 2000 and 28 October 2000). This blazar had thus shown very large peak to peak LTOV during our observations: 1.63 mag within a year.

BL 0735+178: This BL Lac object was observed on four nights between December 1998 and December 2001. Although the data quality was generally good, on no occa-

Figure 2. *Continued*

sion was a clear detection of INOV made, though there is marginal evidence for variations on 24 December 2001 (Fig. 1). This behaviour is in clear contrast to that found here for the remaining five BL Lacs in our sample. Still, substantial LTOV was noted in this BL Lac. It showed significant brightening by 1.15 mag over the course of our first three epochs of observations within two years. The blazar had faded by 0.43 mag from that peak over the next year when last monitored on 24 December 2001.

BL 0851+202: A recent detection of INOV of this well known BL Lac object has been reported by Ghosh et al. (2000). We monitored this object on four nights between

December 1998 and February 2001 and found it to show INOV on all four nights (Fig. 1). On each night features can be seen in the SF, corresponding to time scales between 1.5 and 3.0 hours (Fig. 2). In addition, a “periodicity” of ~ 3.8 hours is present in the SF for the night of 17 February 2001 (Fig. 2; Table 3); however, we note that this corresponds to only two peaks in the light curve (Fig. 1) and so need not be a genuine (quasi-)periodicity. The most remarkable feature on the DLC on 28 March 2000 is a $\sim 4\%$ jump in brightness against all three comparison stars within about an hour (at around 19 UT), which is also borne out in the SF analysis. This blazar also showed very large amplitude

LTOV: it faded by 0.6 mag in a year between 29 December 1998 and 31 December 1999. However it then brightened during our remaining three epochs of observations. A total brightening of 1.63 mag was observed between 31 December 1999 and 17 Feb 2001.

CDQ 0955+326: The only one of the three nights of monitoring this quasar which showed possible indications of variability was 5 March 2000. Against all three comparison stars the quasar showed a steady decline of about 0.5% over 6.9 hours (Fig. 1). Some LTOV of this quasar can be ascertained from our three epochs of observations which cover a span of about one year. The quasar brightened by 0.05 mag within a year between 19 February 1999 and 3 March 2000 and was found to remain at essentially the same level when observed two days later.

CDQ 1128+315: No INOV was detected. Our three epochs of observations span about a year. Between 18 January 2001 and 9 March 2002 the quasar brightened by 0.14 mag; it remained at the same brightness level over the next 24 hours.

BL 1215+303: This BL Lac object was monitored on five nights, of which only four nights were considered for INOV as the data on the remaining one night are of moderate quality. On 20 March 1999 the BL Lac showed clear evidence of variability with peak-to-peak amplitude of ~ 4 to 5% with a 2% change noticed within ~ 0.5 hour. On 19 April 2002 the DLCs showed a steady brightening of 2% over the 6.8 hours of observations (Fig. 1). On 20 March 1999, the SF derived from 7.0 hours of observations shows a steady rise with a turnover corresponding to a time scale of 4.2 hours; no such SF turnover is present on 19 April 2002 (Fig. 2; Table 3). The five epochs of monitoring observations of this BL Lac covered a temporal baseline of over three years (from 20 March 1999 to 19 April 2002). The quasar exhibited LTOV by fading progressively during the first four epochs, becoming fainter by 0.53 mag between 20 March 1999 and 25 April 2001. By 19 April 2002 it had brightened again by 0.11 mag.

CDQ 1216–010: This highly polarized CDQ was monitored on four nights during March 2002 and clear INOV was detected on all the four nights against both comparison stars (Fig. 1). The SF for 16 March 2002 shows a linear rise all the way to the time lag of ~ 8 hours. On each of the earlier three nights a “periodicity” signal of ~ 3 to 4 hours is seen in the SF (Fig. 2; Table 3). This quasar was observed over a time baseline of only 6 days; nonetheless, the source is found to show significant inter-night variability. The quasar brightened by 0.02 mag between the first two epochs (11 March 2002 and 13 March 2002) followed by 0.11 mag fading during the next two days. It once again had brightened (by 0.14 mag over the next 24 hours) when first observed on 16 March 2002; it continued to brighten throughout that night.

CDQ 1225+317: This most distant quasar in our sample did not show any evidence of INOV in our three roughly equally spaced epochs of observations over a span of two years (between 7 March 1999 and 20 April 2001). However, it showed LTOV, wherein a $\sim 2\%$ brightening was noticed in a year between April 2000 and April 2001. One might attribute this lack of detection to a combination of only three nights of observation and the high redshift, which means that we were examining a relatively short period in the rest frame of the CDQ.

BL 1308+326: This BL Lac object was monitored on six epochs between March 1999 and May 2002; however, it was sufficiently bright for the purpose of INOV detection on just the last three epochs, which fall in March, April and May 2002. INOV was detected only on the night of 17 March 2002 showing a gradual fading by about 2% (Fig. 1), as well as possible very short-term variations. The SF shows a plateau corresponding to a time scale of about 1.5 hours (Fig. 2). Large amplitude LTOV was seen. The quasar was relatively faint during the initial three epochs of our observations. A fading of 0.21 mag was found within the first year (23 March 1999 and 26 April 2000) and it was found in the same level when observed a week later on 3 May 2000. However the quasar had brightened by 1.9 mag when observed about 2 years later on 17 March 2002. It faded once again in about a month by 0.44 mag and then brightened by 0.03 mag over 12 days when observed for the last time on 2 May 2002.

CDQ 1309+355: This quasar was observed for three epochs and showed no INOV, although it did exhibit LTOV. Between the first two epochs, separated by about two years, the quasar decreased in brightness by 0.10 mag, and remained at the same brightness level over the next 24 hours.

4.1 Duty cycle (DC) of intra-night optical variability

Duty cycles of INOV were calculated following the definition of Romero et al. (1999). Since most AGNs do not display variability on each night, duty cycles are best estimated not as a fraction of the variable objects found within a given class, but as the ratio of the time over which objects of the class are seen to vary to the total observing time spent on monitoring the objects in the class. It is thus given as

$$DC = 100 \frac{\sum_{i=1}^n N_i (1/\Delta t_i)}{\sum_{i=1}^n (1/\Delta t_i)} \%, \quad (4)$$

where $\Delta t_i = \Delta t_{i,obs}(1+z)^{-1}$ is the duration (corrected for cosmological redshift) of an i^{th} monitoring session of the source out of a total of n sessions for the selected class, and N_i equals 0 or 1, depending on whether the object was non-variable or variable, respectively, during Δt_i . Results for the different classes (CDQ-LP: low polarization CDQs; CDQ-HP: high polarization CDQs; CDQ-ALL: CDQs; BL: BL Lacs) are shown as histograms in Fig. 3. The shaded portions refer to the subsets showing large variability ($\psi > 3\%$), while the blank portions refer to the INOV detections of lower amplitudes; the cases of probable INOV are not included in these histograms.

The results presented here allow for the first time a comparison of the DC of INOV for the three blazar classes, namely: BL Lacs; high-polarization CDQs, CDQ-HP; and low-polarization CDQs, CDQ-LP (although our sample contains just one CDQ-HP, viz., 1216-010 (Table 1), a significant statement about it is possible, since it was monitored on four nights with an average duration of 7.1 hours per night). In fact, the comparisons can now be made for ranges of variability amplitudes, which we take as $\psi < 3\%$ and $\psi > 3\%$. In GSSW03 we presented a similar comparison for RQQs and BL Lac objects monitored in our program. We see that the DC for INOV detection is high both for BL Lacs (61%) and for the CDQ-HP source (100%), but much




Figure 3. Duty cycles for various classes of blazars and for different amplitude ranges: shaded for $\psi > 3\%$; open for $\psi < 3\%$.

lower for the CDQ-LP sources. Thus, there appears to be a close link between the INOV and the polarized component of optical emission (which is commonly attributed to shocks in the relativistic non-thermal jets). While this pattern is highly suggestive, we must caution that a more definitive claim would require that the classification of the blazars into the two polarization classes be based on measurements made simultaneous to the optical flux monitoring and that the number of CDQ-HP sources be increased.

Further, it is interesting to note from Fig. 3 that not only do the BL Lacs and CDQ-HP exhibit high DC of INOV but also the probability of observing large amplitude INOV ($\psi > 3\%$) is high (DC $\simeq 0.5$) and probably very similar to the probability of small amplitude INOV ($\psi < 3\%$), when we note that our lower limit for clear detections of INOV is $\psi \simeq 1\%$.

Any successful model of INOV should be able to explain this rather unexpected behaviour found here for the blazars.

5 CONCLUSIONS

The observations reported here present the first systematic study of the INOV characteristics of the two Doppler beamed AGN classes, CDQs and BL Lacs, although studies of individual sources of these types have been made over the past decade (e.g., Noble et al. 1997). We monitored both classes of blazars in our sample with equally high sensitivities and for comparable time durations; the mean monitoring duration for BL Lacs was 6.5 hours over 31 nights, while for CDQs it was 6.7 hours over 16 nights.

The major result of this observational programme is our finding that the duty cycles of INOV detection for weakly polarized CDQs and BL Lacs are strikingly different. BL Lacs show a high DC of $\sim 60\%$, in contrast to CDQs-LP for which the DC is found to be 0% as none of them showed INOV at the level of 1%. Including the one case of probable variable (0955+326 on 5 March 2000) raises this DC to 6% only. This value is nominally lower than the DCs we have found for RQQs and LDQs ($\sim 15\%$; Stalin et al. 2003a; GSSW03). At the same time, we find a close resem-

blance, both in amplitude and duty cycle of INOV, between the one CDQ-HP in our sample and the BL Lacs. Thus it appears that the mere presence of a prominent (and hence presumably Doppler boosted) radio core does not guarantee INOV; instead, the more crucial factor appears to be the optical polarization of the core emission. Such highly polarized emission is normally associated with shocks in a relativistic jet, so this may not be surprising. Of course the number of sources in our sample is not large, and moreover the polarization measurements were made long before the intra-night optical monitoring reported here. Both these shortcomings need to be overcome in subsequent studies in order to place the present results on a firmer basis.

For most of the sources that do show INOV, some type of time scale of a few hours (observer’s frame) is frequently detected. Although, in a few cases, the structure function indicates the presence of several hour “periodicities”, we stress that these may well correspond to the time gap between multiple flares (or perturbations on bigger, slower flares). None of the light curves are long enough to actually detect putative real periodicities (or quasi-periodicities) of longer than 1 or 2 hours. Probing such phenomena would require continuous monitoring through coordinated observations by many observatories, well separated in longitude, or by a space observatory.

ACKNOWLEDGMENTS

We thank the anonymous referee for suggestions that have significantly improved the presentation of these results. This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. CSS thanks NCRA for hospitality and use of its facilities. PJW is grateful for continuing hospitality at the Department of Astrophysical Sciences at Princeton University; his efforts were partially supported by Research Program Enhancement funds at GSU.

REFERENCES

- Blandford R., Rees M.J., 1978, in Pittsburgh Conference on BL Lac Objects, A. M. Wolfe, ed., U. Pittsburgh, Pittsburgh, p. 328
- Carini M. T., Miller H. R., Noble J. C., Sadun A. C., 1991, AJ, 101, 1196
- Dai B. Z., Xie, G. Z., Li, K. H., Zhou, S. B., Lin, W. W., Jaing, Z. J., 2001, AJ, 122, 2901
- Gabuzda D. C., Cawthorne T. V., Roberts D. H., Wardle J. F. C., 1992, ApJ, 388, 40
- Garcia A., Sodre L., Jablonski F. J., Terlevich R. J., 1999, MNRAS, 309, 803
- Ghosh K. K., Ramsey B. D., Sadun A. C., Soundarajaperumal S., 2000, ApJS, 127, 11
- Gopal-Krishna, Wiita P.J., 1993, Nature, 363, 142
- Gopal-Krishna, Sagar R., Wiita P.J., 1995, MNRAS, 274, 701
- Gopal-Krishna, Stalin C. S., Sagar R., Wiita P. J., 2003, ApJ, 586, L25 (GSSW03)
- Heidt J., Wagner S. J., 1996, A&A, 305, 42
- Hughes P., Aller H., Aller M. 1992, ApJ, 396, 469
- Jang M., Miller H.R., 1997, AJ, 114, 565

- Kovalev Y. Y., Nizhelsky N. A., Kovalev Yu. A., Berlin A. B., Zhekanis G. V., Mingaliev, M. G., Bogdantsov, A. V., 1999, *A&AS*, 139, 545
- Marscher A. P., Gear W. K., 1985, *ApJ*, 298, 114
- Marscher A. P., Gear W. K., Travis J. P., 1992, in Valtaoja E., Valtonen M., eds, *Variability of Blazars*. Cambridge Univ. Press, Cambridge, p. 85
- Mangalam A. V., Wiita P. J., 1993, *ApJ*, 406, 420
- Miller H. R., Carini M. T., Goodrich B. D., 1989, *Nature*, 337, 627
- Noble J. C., Carini M. T., Miller H. R., & Goodrich B., 1997, *AJ*, 113, 1995
- Padovani, P., Giommi, P., 1995, *MNRAS*, 277, 1477
- Romero G. E., Cellone S. A., Combi J. A., 1999, *A&AS*, 135, 477
- Romero G. E., Cellone S. A., Combi J. A., Andruchow I., 2002, *A&A*, 390, 431
- Sagar R., 1999, *Current Science*, 77, 643
- Simonetti J. H., Cordes J. M., Heeschen D. S., 1985, *ApJ*, 296, 46
- Stalin C. S., 2002, Ph.D. dissertation, Kumaun University , Naini Tal
- Stalin C. S., Gopal-Krishna, Sagar R., Wiita P. J., 2003a, accepted in *MNRAS* (astro-ph/0306394)
- Stalin C. S., Gopal-Krishna, Sagar R., Wiita P. J. 2003b, accepted in *JAA* (astro-ph/0306395)
- Stickel, M., Fried, J., Kühr, H., *A&AS*, 98, 393
- Urry C. M., Padovani P., 1995, *PASP*, 107, 803
- Véron-Cetty M.-P., Véron P., 1998, *ESO Scientific Report No. 18*; Publisher: Garching: European Southern Observatory (ESO).
- Wagner S. J., Witzel A., 1995, *ARA&A*, 33, 163
- Wiita P. J., 1996, in Miller H. R., Webb J. R., Noble J. C., eds, *Blazar Continuum Variability*. Astr. Soc. Pacific Conf. Ser. 110, ASP, San Francisco, p. 42
- Wills B. J., Wills D., Breger M., Antonucci R. R. J., Barvainis R., 1992, *ApJ*, 398, 454
- Xie G. Z., Li K. H., Bai J. M., Dai B. Z., Liu W. W., Zhong X., Xing S. Y., 2001, *ApJ*, 548, 200
- Xie G. Z., Zhou S. B., Dai B. Z., Liang E. W., Li K. H., Bai J. M., Xing S. Y., Liu W. W., 2002, *MNRAS*, 329, 689